

UV Spectroscopy of Metal-Poor Massive Stars in the Small Magellanic Cloud

Daniel J. Lennon^{1,2}

¹ Isaac Newton Group of Telescopes djl@ing.iac.es

² Instituto de Astrofísica de Canarias

Summary. The Hubble Space Telescope has provided the first clear evidence for weaker winds of metal-poor massive stars in the Small Magellanic Cloud, confirming theoretical predictions of the metallicity dependence of mass-loss rates and wind terminal velocities. For lower luminosity O-type stars however, derived mass-loss rates are orders of magnitude lower than predicted, and are at present unexplained.

1 Introduction

Massive stars may lose mass throughout their lifetimes either as steady winds or in episodic outbursts during a short-lived Luminous Blue Variable (LBV) phase. To first order, the product of a massive star's mass-loss rate during its core-hydrogen or core-helium burning phase times the duration of these evolutionary phases can amount to a significant fraction of the star's initial mass. Similar considerations apply to the shorter-lived LBV phases. Mass-loss therefore plays a crucial role in the evolution of stars more massive than about 20 solar masses, uncertainties of even a factor of two being significant for the star's evolution. There is general agreement that the steady winds of massive stars are driven by radiation pressure on millions of metal absorption lines, these winds therefore depend on metallicity (Z). Calculations [9] indicate that in the range $0.1Z_{\odot} < Z < 3Z_{\odot}$ the mass-loss rate $\dot{M} \sim Z^x$ where $x \sim 0.7$, while the ratio of wind terminal velocity to escape velocity (v_{∞}/v_{esc}) is rather insensitive to Z , though dependent on effective temperature. The Small Magellanic Cloud with $Z \sim 0.2Z_{\odot}$ is therefore the ideal laboratory in which to study the wind properties and evolution of massive stars at low metallicity.

2 Weak winds in the Small Magellanic Cloud

There were several attempts to confirm theoretical expectations using the IUE satellite to observe massive stars in the Magellanic Clouds, however with mixed results. The Hubble Space Telescope (HST) revolutionized our UV view of massive stars in the Magellanic Clouds, with the first high S/N UV spectroscopic campaign (Kudritzki, G0s:2233/4110) yielding startling




Fig. 1. Spectra of O-type main-sequence stars. Left panel; SMC sources observed with HST/STIS, all of which lie in the cluster NGC 346. Right panel; Galactic sources observed with IUE at high resolution [12]. The features identified at the top are N v $\lambda\lambda$ 1239, 1243; Si iv $\lambda\lambda$ 1394, 1403; and C iv $\lambda\lambda$ 1548, 1551. A characteristic of O-type dwarfs in the SMC is the absence of saturated C iv P-Cygni profiles, in sharp contrast to the Galactic stars. In addition wind features are extremely weak or absent for dwarfs later than O5. The strong N v feature in MPG 355 reflects the large nitrogen enrichment in this object.

spectra from the Faint Object Spectrograph (FOS), whose morphology confirmed the presence of weak winds in the O-type main-sequence stars in the SMC [14]. Subsequent quantitative analysis confirmed that mass-loss rates were indeed lower in the SMC sample than in LMC and Galactic counterparts [8], and in reasonable agreement with theory. This prompted a more systematic approach to the SMC, enlarging the previous sample of 9 stars with an additional 30, the use of high resolution with STIS, the inclusion of B-type supergiants, and an extension to lower luminosity late-O dwarfs (Lennon, GOs:7437/9116; [15], [2]). The UV spectra for O-type dwarfs and B-type supergiants are displayed and discussed in Fig.1 and Fig.2 respectively.

Many of these stars have been analysed to derive mass-loss rates confirming the expected theoretical trend of mass-loss with metallicity ([1], [5], [10], [11], [4]) while theoretical expectations for terminal velocities have also been confirmed [3]. However, a small number of stars were found to have mass-loss rates which are an order of magnitude or more less than theoretical predictions [1], see Fig.3. Additional members of this group have been found in the




Fig. 2. Spectra of B-type supergiant stars. Left panel; SMC sources observed with HST/STIS. Right panel; Galactic sources observed with IUE at high resolution [13]. The features identified at the top are N V $\lambda\lambda$ 1239, 1243; C II $\lambda\lambda$ 1334, 1336; Si IV $\lambda\lambda$ 1394, 1403; C IV $\lambda\lambda$ 1548, 1551; and Al III $\lambda\lambda$ 1855, 1863. Note the extreme weakness of these features in the SMC stars compared to their Galactic counterparts.

SMC [6], in all cases these stars are low luminosity O-type dwarfs having anomalous weak winds. The nature of their winds is not as yet understood, but possibly indicate an incomplete understanding of mass-loss in the weak wind limit, or perhaps arise as a result of the lack of accurate diagnostics.

References

1. Bouret, J.-C., Lanz, T., Hillier, D.J., Heap, S.R., Hubeny, I., Lennon, D.J., Smith, L.J., & Evans, C.J. 2003, *ApJ*, 595, 1182
2. Evans, C.J., Lennon, D.J., Walborn, N.R., Trundle, C. & Rix, S.A. 2004, *PASP*, 116, 909
3. Evans, C.J., Lennon, D.J., Trundle, C., Heap, S.R., & Lindler, D.J. 2004, *ApJ*, 607, 451
4. Evans, C.J., Crowther, P.A., Fullerton, A.W. & Hillier, D.J. 2004, *ApJ*, 610, 1021
5. Hillier, D.J., Lanz, T., Heap, S.R., Hubeny, I., Smith, L.J., Evans, C.J., Lennon, D.J., & Bouret, J.-C. 2003, *ApJ*, 588, 1039
6. Martins, F., Schaerer, D., Hillier, D.J. & Heydari-Malayeri, M. 2004, *A&A*, 420, 1087




Fig. 3. Modified wind-momentum ($D = \dot{M}v_{\infty}\sqrt{R}$) versus luminosity for SMC OB stars observed with HST ([1], [5], [10], [11]); filled symbols represent stars with measured terminal velocities, open symbols are stars with assumed terminal velocities, circles and squares represent O- and B-type stars respectively. Additional OB supergiants from [4] are marked with asterisks. In all cases we refer to unclumped mass-loss rates. The lines are the empirical fits derived for O-type stars [7] (dashed line), early B-type supergiants [11] (dash-dot) and mid B-type supergiants [11] (dash-dot-dot-dot). The three O-type stars (open circles) well below the expected relationship, plus the upper limits (inverted triangles) from [6] represent the anomalous weak wind class of O-type stars.

7. Mokiem, M.R., de Koter, A., Vink, J.S., Puls, J., Evans, C.J., Smartt, S.J., Crowther, P.A., Herrero, A., Langer, N., Lennon, D.J., Najarro, F. & Villamariz, M.R. 2007, A&A, in press.
8. Puls, J., Kudritzki, R.-P., Herrero, A., Pauldrach, A.W.A., Haser, S.M., Lennon, D.J., Gabler, R., Voels, S.A., Vilchez, J.M., Wachter, S. & Feldmeier, A. 1996, A&A, 305, 171
9. Puls, J., Springmann, U., & Lennon, M.A. 2000, A&AS, 141, 23
10. Trundle, C., Lennon, D.J., Puls, J., & Dufton, P.L. 2004, A&A, 417, 217
11. Trundle, C. & Lennon, D.J. 2005, A&A, 434, 677
12. Walborn, N.R., Nichols-Bohlin, J. & Panek, R.J.. 1985, International Ultraviolet Explorer Atlas of O-type Spectra (NASA RP1155; Washington; NASA)
13. Walborn, N.R., Parker, J.W., & Nichols, J. 1995, International Ultraviolet Explorer Atlas of B-type Spectra (NASA RP1363; Washington; NASA)
14. Walborn, N.R., Lennon, D.J., Haser, S.M., Kudritzki, R.-P., & Voels, S.A. 1995, PASP, 107, 104
15. Walborn, N.R., Lennon, D.J., Heap, S.R., Lindler, D.J., Smith, L., Evans, C.J. & Parker, J.W. 2000, PASP, 112, 1243